

Calculate derivatives of each of the trigonometric ratios, given that you know  $\frac{d}{dx}[\sin(x)] = \cos(x)$  and  $\frac{d}{dx}[\cos(x)] = -\sin(x)$ .

Sol: We'll calculate derivatives of ①  $\tan(x)$ , ②  $\sec(x)$ , ③  $\csc(x)$ , and ④  $\cot(x)$ .

$$\begin{aligned}\textcircled{A} \quad \frac{d}{dx}[\tan(x)] &= \frac{d}{dx}\left[\frac{\sin(x)}{\cos(x)}\right] = \frac{\cos(x)\frac{d}{dx}[\sin(x)] - \sin(x)\frac{d}{dx}[\cos(x)]}{(\cos(x))^2} \\ &= \frac{\cos(x) \cdot \cos(x) - \sin(x) \cdot (-\sin(x))}{\cos^2(x)} \\ &= \frac{\cos^2(x) + \sin^2(x)}{\cos^2(x)} = \frac{1}{\cos^2(x)} = \sec^2(x)\end{aligned}$$

$$\begin{aligned}\textcircled{B} \quad \frac{d}{dx}[\sec(x)] &= \frac{d}{dx}\left[\frac{1}{\cos(x)}\right] = \frac{\cos(x)\frac{d}{dx}[1] - 1 \cdot \frac{d}{dx}[\cos(x)]}{(\cos(x))^2} \\ &= \frac{\cos(x) \cdot 0 - (-\sin(x))}{\cos^2(x)} \\ &= \frac{\sin(x)}{\cos^2(x)} = \frac{1}{\cos(x)} \cdot \frac{\sin(x)}{\cos(x)} = \sec(x) \tan(x)\end{aligned}$$

$$\begin{aligned}\textcircled{C} \quad \frac{d}{dx}[\csc(x)] &= \frac{d}{dx}\left[\frac{1}{\sin(x)}\right] = \frac{\sin(x) \cdot \frac{d}{dx}[1] - 1 \cdot \frac{d}{dx}[\sin(x)]}{(\sin(x))^2} \\ &= \frac{\sin(x) \cdot 0 - \cos(x)}{\sin^2(x)} \\ &= -\frac{\cos(x)}{\sin^2(x)} = -\frac{1}{\sin(x)} \cdot \frac{\cos(x)}{\sin(x)} = -\csc(x) \cot(x)\end{aligned}$$

$$\begin{aligned}\textcircled{D} \quad \frac{d}{dx}[\cot(x)] &= \frac{d}{dx}\left[\frac{\cos(x)}{\sin(x)}\right] = \frac{\sin(x)\frac{d}{dx}[\cos(x)] - \cos(x)\frac{d}{dx}[\sin(x)]}{(\sin(x))^2} \\ &= \frac{\sin(x) \cdot (-\sin(x)) - \cos(x) \cdot \cos(x)}{\sin^2(x)} \\ &= -\left(\frac{\sin^2(x) + \cos^2(x)}{\sin^2(x)}\right) \\ &= -\frac{1}{\sin^2(x)} = -\csc^2(x)\end{aligned}$$

# DERIVATIVES OF TRIGONOMETRIC FUNCTIONS

175-184: Calculate  $\frac{dy}{dx}$ .

175.  $y = x^2 - \sec(x) + 1$

Sol:  $\frac{dy}{dx} = \frac{d}{dx}[x^2 - \sec(x) + 1] = 2x - \sec(x)\tan(x)$   $\boxed{\text{G}}$

176.  $y = 3\csc(x) + \frac{5}{x}$

Sol:  $\frac{dy}{dx} = \frac{d}{dx}[3\csc(x) + 5x^{-1}] = -3\csc(x)\cot(x) - 5x^{-2}$   $\boxed{\text{G}}$

177.  $y = x^2 \cot(x)$

Sol:  $\frac{dy}{dx} = \frac{d}{dx}[x^2 \cot(x)] = \frac{d}{dx}[x^2]\cot(x) + x^2 \frac{d}{dx}[\cot(x)]$   
 $= 2x \cot(x) - x^2 \csc^2(x)$   $\boxed{\text{G}}$

178.  $y = x - x^3 \sin(x)$

Sol:  $\frac{dy}{dx} = \frac{d}{dx}[x - x^3 \sin(x)] = 1 - \left( \frac{d}{dx}[x^3] \sin(x) + x^3 \frac{d}{dx}[\sin(x)] \right)$   
 $= 1 - 3x^2 \sin(x) - x^3 \cos(x)$   $\boxed{\text{G}}$

179.  $y = \frac{\sec(x)}{x}$

Sol:  $\frac{dy}{dx} = \frac{d}{dx}\left[\frac{\sec(x)}{x}\right] = \frac{x \frac{d}{dx}[\sec(x)] - \sec(x) \frac{d}{dx}[x]}{x^2}$   
 $= \frac{x \sec(x) \tan(x) - \sec(x)}{x^2}$   $\boxed{\text{G}}$

$$180. \quad y = \sin(x) \tan(x)$$

$$\begin{aligned} \text{Sol: } \frac{dy}{dx} &= \frac{d}{dx} [\sin(x) \tan(x)] = \frac{d}{dx} [\sin(x)] \tan(x) + \sin(x) \frac{d}{dx} [\tan(x)] \\ &= \cos(x) \tan(x) + \sin(x) \cdot \sec^2(x) \\ &= \cos(x) \cdot \frac{\sin(x)}{\cos(x)} + \sin(x) \sec^2(x) \\ &= \sin(x) (1 + \sec^2(x)) \quad \square \end{aligned}$$

$$181. \quad y = (x + \cos(x))(1 - \sin(x))$$

$$\begin{aligned} \text{Sol: } \frac{dy}{dx} &= \frac{d}{dx} [(x + \cos(x))(1 - \sin(x))] \\ &= \frac{d}{dx} [x + \cos(x)] (1 - \sin(x)) + (x + \cos(x)) \frac{d}{dx} [1 - \sin(x)] \\ &= (1 - \sin(x))(1 - \sin(x)) + (x + \cos(x)) \cdot -\cos(x) \\ &= 1 - 2\sin(x) + \sin^2(x) - x\cos(x) - \cos^2(x) \quad \square \end{aligned}$$

$$182. \quad y = \frac{\tan(x)}{1 - \sec(x)}$$

$$\begin{aligned} \text{Sol 1: } \frac{dy}{dx} &= \frac{d}{dx} \left[ \frac{\tan(x)}{1 - \sec(x)} \right] = \frac{(1 - \sec(x)) \frac{d}{dx} [\tan(x)] - \tan(x) \frac{d}{dx} [1 - \sec(x)]}{(1 - \sec(x))^2} \\ &= \frac{(1 - \sec(x)) \sec^2(x) - \tan(x) \cdot -\sec(x) \tan(x)}{(1 - \sec(x))^2} \\ &= \frac{\sec(x) (\sec(x) - \sec^2(x) + \tan^2(x))}{(1 - \sec(x))^2} \quad \square \end{aligned}$$

$$\begin{aligned} \text{Sol 2: } \frac{dy}{dx} &= \frac{d}{dx} \left[ \frac{\tan(x)}{1 - \sec(x)} \right] = \frac{d}{dx} \left[ \frac{\sin(x)/\cos(x)}{1 - \cos(x)} \cdot \frac{\cos(x)}{\cos(x)} \right] - \frac{d}{dx} \left[ \frac{\sin(x)}{\cos(x) - 1} \right] \\ &= \frac{(\cos(x) - 1) \frac{d}{dx} [\sin(x)] - \sin(x) \frac{d}{dx} [\cos(x) - 1]}{(\cos(x) - 1)^2} \\ &= \frac{(\cos(x) - 1) \cos(x) - \sin(x) \cdot -\sin(x)}{(\cos(x) - 1)^2} = \frac{\cos^2(x) - \cos(x) + \sin^2(x)}{(\cos(x) - 1)^2} \\ &= \frac{1 - \cos(x)}{(\cos(x) - 1)^2} = \frac{1}{1 - \cos(x)} \quad \square \end{aligned}$$

$$183. \quad y = \frac{1 - \cot(x)}{1 + \cot(x)}$$

$$\begin{aligned} \underline{\text{So 1:}} \quad \frac{dy}{dx} &= \frac{d}{dx} \left[ \frac{1 - \cot(x)}{1 + \cot(x)} \right] = \frac{(1 + \cot(x)) \frac{d}{dx}[1 - \cot(x)] - (1 - \cot(x)) \frac{d}{dx}[1 + \cot(x)]}{(1 + \cot(x))^2} \\ &= \frac{(1 + \cot(x)) \cdot -(-\operatorname{sc}^2(x)) - (1 - \cot(x)) \cdot -\operatorname{sc}^2(x)}{(1 + \cot(x))^2} \\ &= \frac{\operatorname{sc}^2(x) (1 + \cot(x) - 1 + \cot(x))}{(1 + \cot(x))^2} = \frac{2 \operatorname{sc}^2(x) \cot(x)}{(1 + \cot(x))^2} \quad \boxed{B} \end{aligned}$$

$$\begin{aligned} \underline{\text{So 2:}} \quad \frac{dy}{dx} &= \frac{d}{dx} \left[ \frac{1 - \cot(x)}{1 + \cot(x)} \right] = \frac{d}{dx} \left[ \frac{1 - \frac{\cos(x)}{\sin(x)}}{1 + \frac{\cos(x)}{\sin(x)}} \cdot \frac{\sin(x)}{\sin(x)} \right] = \frac{d}{dx} \left[ \frac{\sin(x) - \cos(x)}{\sin(x) + \cos(x)} \right] \\ &= \frac{(\sin(x) + \cos(x)) \frac{d}{dx}[\sin(x) - \cos(x)] - (\sin(x) - \cos(x)) \frac{d}{dx}[\sin(x) + \cos(x)]}{(\sin(x) + \cos(x))^2} \\ &= \frac{(\sin(x) + \cos(x))(\cos(x) + \sin(x)) - (\sin(x) - \cos(x))(\cos(x) - \sin(x))}{(\sin(x) + \cos(x))^2} \\ &= \frac{(\sin(x) + \cos(x))^2 + (\sin(x) - \cos(x))^2}{(\sin(x) + \cos(x))^2} \\ &= \frac{\sin^2(x) + 2\sin(x)\cos(x) + \cos^2(x) + \sin^2(x) - 2\sin(x)\cos(x) + \cos^2(x)}{(\sin(x) + \cos(x))^2} \\ &= \frac{2(\sin^2(x) + \cos^2(x))}{(\sin(x) + \cos(x))^2} = \frac{2}{(\sin(x) + \cos(x))^2} \quad \boxed{VII} \end{aligned}$$

$$184. \quad y = \cos(x) (1 + \operatorname{sc}(x))$$

$$\begin{aligned} \underline{\text{So 1:}} \quad \frac{dy}{dx} &= \frac{d}{dx} [\cos(x) (1 + \operatorname{sc}(x))] = \frac{d}{dx} [\cos(x)] (1 + \operatorname{sc}(x)) + \cos(x) \frac{d}{dx} [1 + \operatorname{sc}(x)] \\ &= -\sin(x) \cdot (1 + \operatorname{sc}(x)) + \cos(x) (-\operatorname{sc}(x) \cot(x)) \leftarrow \text{OK} \\ &= -\sin(x) \left( 1 + \frac{1}{\sin(x)} \right) - \cos(x) \left( \frac{1}{\sin(x)} \cdot \frac{\cos(x)}{\sin(x)} \right) \\ &= -\sin(x) - (1 + \cot^2(x)) \quad \leftarrow 1 + \cot^2(x) = \operatorname{sc}^2(x) \\ &= -\sin(x) - \operatorname{sc}^2(x) \quad \boxed{G} \end{aligned}$$

185-190: Find the equation of the tangent line to  $y=f(x)$  at the given value of  $x$ .

Recall: ①  $f'(a)$  is the slope of the tangent line at  $a$ .  
 ② Lines have point-slope form  $y-y_0 = m(x-x_0)$ .  
 ③ I give my answers in both point-slope form and slope intercept form, but you can choose to only give one (usually point-slope is easier).

185.  $f(x) = -\sin(x)$  at  $x=0$

Sol:  $f'(x) = \frac{d}{dx}[-\sin(x)] = -\frac{d}{dx}[\sin(x)] = -\cos(x)$ .

$$\therefore m = f'(0) = -\cos(0) = -1$$

$$f(0) = -\sin(0) = -0 = 0$$

$$\therefore \text{tangent line is } y - 0 = -1(x - 0) \text{ or } y = -x. \quad \boxed{\checkmark}$$

186.  $f(x) = \csc(x)$  at  $x=\frac{\pi}{2}$

Sol:  $f'(x) = \frac{d}{dx}[\csc(x)] = -\csc(x)\cot(x)$

$$\therefore m = f'\left(\frac{\pi}{2}\right) = -\csc\left(\frac{\pi}{2}\right)\cot\left(\frac{\pi}{2}\right) = -\frac{1}{\sin\left(\frac{\pi}{2}\right)} \cdot \frac{\cos\left(\frac{\pi}{2}\right)}{\sin\left(\frac{\pi}{2}\right)} = -\frac{1}{1} \cdot \frac{0}{1} = 0$$

$$f\left(\frac{\pi}{2}\right) = \csc\left(\frac{\pi}{2}\right) = \frac{1}{\sin\left(\frac{\pi}{2}\right)} = \frac{1}{1} = 1$$

$$\therefore y - 1 = 0(x - \frac{\pi}{2}) \quad \text{OR} \quad y = 1 \quad \boxed{\checkmark}$$

187.  $f(x) = 1 + \cos(x)$  at  $x=\frac{3\pi}{2}$

Sol:  $f'(x) = \frac{d}{dx}[1 + \cos(x)] = -\sin(x)$

$$m = f'\left(\frac{3\pi}{2}\right) = -\sin\left(\frac{3\pi}{2}\right) = -(-1) = 1$$

$$f\left(\frac{3\pi}{2}\right) = 1 + \cos\left(\frac{3\pi}{2}\right) = 1 + 0 = 1$$

$$\therefore y - 1 = 1(x - \frac{3\pi}{2}) \quad \text{OR} \quad y = x + 1 - \frac{3\pi}{2} \quad \boxed{\checkmark}$$

$$\underline{188} \quad f(x) = \sec(x) \quad \text{at} \quad x = \frac{\pi}{4}$$

$$\text{Sol: } f'(x) = \sec(x) \tan(x)$$

$$m = f'\left(\frac{\pi}{4}\right) = \sec\left(\frac{\pi}{4}\right) \tan\left(\frac{\pi}{4}\right) = \frac{1}{\cos\left(\frac{\pi}{4}\right)} \cdot \frac{\sin\left(\frac{\pi}{4}\right)}{\cos\left(\frac{\pi}{4}\right)} = \frac{\sqrt{2}/2}{(\sqrt{2}/2)(\sqrt{2}/2)} = \frac{2}{\sqrt{2}} = \sqrt{2}$$

$$f\left(\frac{\pi}{4}\right) = \sec\left(\frac{\pi}{4}\right) = \frac{1}{\cos\left(\frac{\pi}{4}\right)} = \frac{1}{\sqrt{2}/2} = \frac{2}{\sqrt{2}} = \sqrt{2}$$

$$\therefore y - \sqrt{2} = \sqrt{2}(x - \frac{\pi}{4}) \quad \text{OR} \quad y = \sqrt{2}x + \sqrt{2}\left(1 - \frac{\pi}{4}\right) \quad \boxed{\checkmark}$$

$$\underline{189} \quad f(x) = x^2 - \tan(x) \quad \text{at} \quad x = 0$$

$$\text{Sol: } f'(x) = 2x - \sec^2(x)$$

$$m = f'(0) = 2 \cdot 0 - \sec^2(0) = -\frac{1}{\cos^2(0)} = -\frac{1}{1^2} = -1$$

$$f(0) = 0^2 - \tan(0) = 0 - 0 = 0$$

$$\therefore y - 0 = -1(x - 0) \quad \text{OR} \quad y = -x \quad \boxed{\checkmark}$$

$$\underline{190} \quad f(x) = 5 \cot(x) \quad \text{at} \quad x = \frac{\pi}{4}$$

$$\text{Sol: } f'(x) = -5 \csc^2(x)$$

$$m = -5 \csc^2\left(\frac{\pi}{4}\right) = -5 \cdot \frac{1}{\sin^2\left(\frac{\pi}{4}\right)} = -5 / (\sqrt{2}/2)^2 = -5 / (2/4) = -10$$

$$f\left(\frac{\pi}{4}\right) = 5 \cot\left(\frac{\pi}{4}\right) = 5 \cdot \frac{\cos\left(\frac{\pi}{4}\right)}{\sin\left(\frac{\pi}{4}\right)} = 5 \cdot \frac{\sqrt{2}/2}{\sqrt{2}/2} = 5 \cdot 1 = 5$$

$$\therefore y - 5 = -10(x - \frac{\pi}{4}) \quad \text{OR} \quad y = -10x + 5\left(\frac{\pi}{2} + 1\right)$$

191 - 196: Calculate  $\frac{d^2y}{dx^2}$  for the given  $y = f(x)$ .

Recall:  $\frac{d^2y}{dx^2} = \frac{d}{dx} \left[ \frac{dy}{dx}[y] \right]$  is the second derivative.

191.  $y = x \sin(x) - \cos(x)$

Sol:  $\frac{dy}{dx} = \frac{d}{dx} [x \sin(x) - \cos(x)]$   
=  $x \cos(x) + 1 \cdot \sin(x) - (-\sin(x))$   
=  $x \cos(x) + 2 \sin(x)$

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx} [x \cos(x) + 2 \sin(x)] \\&= x(-\sin(x)) + 1 \cdot \cos(x) + 2 \cos(x) \\&= 3 \cos(x) - x \sin(x) \quad \boxed{\text{II}}\end{aligned}$$

192.  $y = \sin(x) \cos(x)$

Sol:  $\frac{dy}{dx} = \frac{d}{dx} [\sin(x) \cos(x)]$   
=  $\cos(x) \cdot \cos(x) + \sin(x) \cdot (-\sin(x))$   
=  $\cos^2(x) - \sin^2(x)$

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx} [\cos^2(x) - \sin^2(x)] \quad \downarrow \text{generalized power rule, twice} \\&= 2 \cos(x) \cdot (-\sin(x)) - 2 \sin(x) \cdot \cos(x) \\&= -4 \sin(x) \cos(x) \quad \boxed{\text{III}}\end{aligned}$$

193.  $y = x - \frac{1}{2} \sin(x)$

Sol:  $\frac{dy}{dx} = \frac{d}{dx} \left[ x - \frac{1}{2} \sin(x) \right] = 1 - \frac{1}{2} \cdot \cos(x)$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} \left[ 1 - \frac{1}{2} \cos(x) \right] = 0 - \frac{1}{2} (-\sin(x)) = \frac{1}{2} \sin(x) \quad \boxed{\text{IV}}$$

$$\underline{194.} \quad y = \frac{1}{x} + \tan(x) = x^{-1} + \tan(x)$$

$$\underline{\text{Sol:}} \quad \frac{dy}{dx} = \frac{d}{dx} \left[ x^{-1} + \tan(x) \right] = -x^{-2} + \sec^2(x)$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{d}{dx} \left[ -x^{-2} + \sec^2(x) \right] = -(-2x^{-3}) + 2\sec(x) \cdot \sec(x) \tan(x) \\ &= 2x^{-3} + 2\sec^2(x) \tan(x) \quad \boxed{16} \end{aligned}$$

$$\underline{195.} \quad y = 2\csc(x)$$

$$\underline{\text{Sol:}} \quad \frac{dy}{dx} = \frac{d}{dx} [2\csc(x)] = 2 \cdot -(\csc(x) \cot(x)) = -2\csc(x) \cot(x).$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{d}{dx} [-2\csc(x) \cot(x)] = -2 \left( (\csc(x) \frac{d}{dx} [\cot(x)] + \frac{d}{dx} [\csc(x)] \cot(x)) \right) \\ &= -2 \left( \csc(x) \cdot -\csc^2(x) + (-\csc(x) \cot(x)) \cot(x) \right) \\ &= 2\csc(x) (\csc^2(x) + \cot^2(x)) \quad \boxed{16} \end{aligned}$$

$$\underline{196.} \quad y = \sec^2(x)$$

$$\underline{\text{Sol:}} \quad \frac{dy}{dx} = \frac{d}{dx} [\sec^2(x)] = 2\sec(x) \cdot \sec(x) \tan(x) = 2\sec^2(x) \tan(x)$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{d}{dx} [2\sec^2(x) \tan(x)] = 2 \left( \sec^2(x) \frac{d}{dx} [\tan(x)] + \frac{d}{dx} [\sec^2(x)] \tan(x) \right) \\ &= 2 \left( \sec^2(x) \cdot \sec^2(x) + 2\sec^2(x) \tan(x) \cdot \tan(x) \right) \\ &= 2\sec^2(x) (\sec^2(x) + 2\tan^2(x)) \quad \boxed{16} \end{aligned}$$

200. A mass on a spring bounces up and down in simple harmonic motion, modeled by the function  $s(t) = -6\cos(t)$  where  $s$  is measured in inches and  $t$  is measured in seconds.

Calculate the rate at which the spring is oscillating after 5 seconds.

Sol: The desired quantity is  $s'(5)$

$$s'(t) = \frac{d}{dt}[-6\cos(t)] = -6 \cdot -\sin(t) = 6\sin(t).$$

$$s'(5) = 6\sin(5) \leftarrow \text{exact value!}$$

∴ the spring oscillates at a rate of  $6\sin(5)$  inches/sec.  $\boxed{\boxed{}}$

201. Let the position of a swinging pendulum in simple harmonic motion be given by  $s(t) = a\cos(t) + b\sin(t)$  where  $a$  and  $b$  are constants,  $t$  measures time in seconds, and  $s$  measures position in centimeters. If the position is 0 cm and the velocity is 3 cm/s when  $t=0$ , calculate  $a$  and  $b$ .

Sol: We are told:  $\begin{cases} s(0) = 0 \\ s'(0) = 3 \end{cases}$

$$\therefore 0 = s(0) = a\cos(0) + b\sin(0) = a \cdot 1 + b \cdot 0 = a$$

$$\begin{aligned} s'(t) &= \frac{d}{dt}[a\cos(t) + b\sin(t)] \\ &= a(-\sin(t)) + b\cos(t) \\ &= b\cos(t) - a\sin(t) \end{aligned}$$

$$3 = s'(0) = b\cos(0) - a\sin(0) = b \cdot 1 - a \cdot 0 = b$$

$$\therefore a = 0 \quad \text{and} \quad b = 3 \quad \boxed{\boxed{}}$$

209-213. Calculate the given higher-order derivative.

209.  $\frac{d^3y}{dx^3}$  for  $y = 3 \cos(x)$ .

Sol:  $\frac{dy}{dx} = -3 \sin(x)$ ,  $\frac{d^2y}{dx^2} = \frac{d}{dx}[-3 \sin(x)] = -3 \cos(x)$

$$\frac{d^3y}{dx^3} = \frac{d}{dx}[-3 \cos(x)] = 3 \sin(x) \quad \boxed{11}$$

210.  $\frac{d^2y}{dx^2}$  for  $y = 3 \sin(x) + x^2 \cos(x)$

Sol:  $\frac{dy}{dx} = \frac{d}{dx}[3 \sin(x) + x^2 \cos(x)]$   
 $= \frac{d}{dx}[3 \sin(x)] + \frac{d}{dx}[x^2 \cos(x)]$   
 $= 3 \cos(x) + \frac{d}{dx}[x^2] \cos(x) + x^2 \frac{d}{dx}[\cos(x)]$   
 $= 3 \cos(x) + 2x \cos(x) + x^2(-\sin(x))$   
 $= (2x+3) \cos(x) - x^2 \sin(x)$

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx}[(2x+3) \cos(x) - x^2 \sin(x)] \\&= \frac{d}{dx}[(2x+3) \cos(x)] - \frac{d}{dx}[x^2 \sin(x)] \\&= \left( \frac{d}{dx}[2x+3] \cos(x) + \frac{d}{dx}[\cos(x)](2x+3) \right) \\&\quad - \left( \frac{d}{dx}[x^2] \sin(x) + \frac{d}{dx}[\sin(x)] x^2 \right) \\&= (2 \cos(x) + (-\sin(x))(2x+3)) - (2x \sin(x) + \cos(x) \cdot x^2) \\&= 2 \cos(x) - (2x+3) \sin(x) - 2x \sin(x) + x^2 \cos(x) \\&= (x^2+2) \cos(x) - (4x+3) \sin(x) \quad \boxed{12}\end{aligned}$$

$$211. \frac{d^4y}{dx^4} \text{ for } y = 5 \cos(x)$$

$$\text{Sol: } \frac{dy}{dx} = -5 \sin(x), \quad \frac{d^2y}{dx^2} = -5 \cos(x)$$

$$\frac{d^3y}{dx^3} = 5 \sin(x), \quad \frac{d^4y}{dx^4} = 5 \cos(x) \quad \square$$

$$212. \frac{d^2y}{dx^2} \text{ for } y = \sec(x) + \cot(x)$$

$$\text{Sol: } \frac{dy}{dx} = \frac{d}{dx} [\sec(x) + \cot(x)] = \sec(x)\tan(x) - \csc^2(x)$$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} [\sec(x)\tan(x) - \csc^2(x)]$$

$$= \frac{d}{dx} [\sec(x)\tan(x)] - \frac{d}{dx} [\csc(x) \cdot \csc(x)]$$

$$= \left( \frac{d}{dx} [\sec(x)] \tan(x) + \frac{d}{dx} [\tan(x)] \sec(x) \right)$$

$$- \left( \frac{d}{dx} [\csc(x)] \csc(x) + \frac{d}{dx} [\csc(x)] \csc(x) \right)$$

$$= \sec(x)\tan(x) \cdot \tan(x) + \sec^2(x) \cdot \sec(x) - 2(-\csc(x)\cot(x))(\csc(x))$$

$$= \sec(x)\tan^2(x) + \sec^3(x) + 2\csc^2(x)\cot(x) \quad \square$$

$$\underline{213.} \quad \frac{d^3y}{dx^3} \quad \text{for } y = x^{10} - \sec(x)$$

$$\underline{\text{Sol}}: \quad \frac{dy}{dx} = \frac{d}{dx} [x^{10} - \sec(x)] = 10x^9 - \sec(x)\tan(x)$$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} [10x^9 - \sec(x)\tan(x)] = \frac{d}{dx} [10x^9] - \frac{d}{dx} [\sec(x)\tan(x)]$$

$$= 90x^8 - \left( \frac{d}{dx} [\sec(x)] \tan(x) + \sec(x) \frac{d}{dx} [\tan(x)] \right)$$

$$= 90x^8 - (\sec(x)\tan(x)\tan(x) + \sec(x)\cdot\sec^2(x))$$

$$= 90x^8 - \sec(x)(\tan^2(x) + \sec^2(x))$$

$$\frac{d^3y}{dx^3} = \frac{d}{dx} [90x^8 - \sec(x)(\tan^2(x) + \sec^2(x))]$$

$$= \frac{d}{dx} [90x^8] - \frac{d}{dx} [\sec(x)(\tan^2(x) + \sec^2(x))]$$

$$= 720x^7 - \left( \frac{d}{dx} [\sec(x)] (\tan^2(x) + \sec^2(x)) + \sec(x) \frac{d}{dx} [\tan^2(x) + \sec^2(x)] \right)$$

$$= 720x^7 - (\sec(x)\tan(x)(\tan^2(x) + \sec^2(x))$$

$$+ \sec(x)(2\tan(x)\sec^2(x) + 2\sec(x)\cdot\sec(x)\tan(x)))$$

$$= 720x^7 - \sec(x)(\tan^3(x) + \sec^2(x)\tan(x) + 4\sec^2(x)\tan(x))$$

$$= 720x^7 - \sec(x)\tan(x)(\tan^2(x) + 5\sec^2(x)) \quad \boxed{4}$$

Note: It's possible your answer looks different but is still correct (if you used any trig identities, for example).