

DERIVATIVE PRACTICE

1. Calculate the derivative $f'(x)$ for...

a. $f(x) = \frac{x^3 - 6x}{1 + 2x^2}$

Sol: $f'(x) = \frac{d}{dx} \left[\frac{x^3 - 6x}{1 + 2x^2} \right] = \frac{(1 + 2x^2) \frac{d}{dx} [x^3 - 6x] - (x^3 - 6x) \frac{d}{dx} [1 + 2x^2]}{(1 + 2x^2)^2}$

$$= \frac{(1 + 2x^2)(3x^2 - 6) - (x^3 - 6x)(4x)}{(1 + 2x^2)^2}$$
$$= \frac{3x^2 - 6 + 6x^4 - 12x^2 - 4x^4 + 24x^2}{(1 + 2x^2)^2}$$
$$= \frac{2x^4 + 15x^2 - 6}{(1 + 2x^2)^2} \quad \square$$

b. $f(x) = x^2 \cos(x) - x \sin(x)$

Sol: $f'(x) = \frac{d}{dx} [x^2 \cos(x) - x \sin(x)] = \frac{d}{dx} [x^2 \cos(x)] - \frac{d}{dx} [x \sin(x)]$

$$= \left(\frac{d}{dx} [x^2] \cos(x) + x^2 \frac{d}{dx} [\cos(x)] \right) - \left(\frac{d}{dx} [x] \sin(x) + x \frac{d}{dx} [\sin(x)] \right)$$
$$= 2x \cos(x) + x^2 (-\sin(x)) - \sin(x) - x \cos(x)$$
$$= x \cos(x) - (x^2 + 1) \sin(x) \quad \square$$

c. $f(x) = \frac{x^2 - 1}{\cos(x)}$

Sol: $f'(x) = \frac{d}{dx} \left[\frac{x^2 - 1}{\cos(x)} \right] = \frac{\cos(x) \frac{d}{dx} [x^2 - 1] - (x^2 - 1) \frac{d}{dx} [\cos(x)]}{\cos^2(x)}$

$$= \frac{\cos(x) \cdot 2x - (x^2 - 1)(-\sin(x))}{\cos^2(x)}$$
$$= \frac{2x \cos(x) + (x^2 - 1) \sin(x)}{\cos^2(x)} \quad \square$$

2. Find an equation to the tangent line to $y=f(x)$ at $x=a$.

2a. $y = x^2 + \frac{15}{x} - 10$, $a = 5$

Sol: $y - y_0 = m(x - x_0)$ ← equation of a line

$$y'(x) = 2x - 15x^{-2} = 2x - \frac{15}{x^2}$$

$$m = y'(5) = 2 \cdot 5 - \frac{15}{5^2} = 10 - \frac{3}{5} = \frac{47}{5}$$

$$x_0 = 5, \quad y_0 = y(5) = 5^2 + \frac{15}{5} - 10 = 18$$

$$y - 18 = \frac{47}{5}(x - 5) \quad \square$$

2b. $y = (3x - x^2)(3 - x - x^2)$, $a = 1$

Sol: $y - y_0 = m(x - x_0)$ ← equation of a line

$$y'(x) = \frac{d}{dx}[3x - x^2](3 - x - x^2) + (3x - x^2) \frac{d}{dx}[3 - x - x^2]$$

$$= (3 - 2x)(3 - x - x^2) + (3x - x^2)(-1 - 2x)$$

$$m = y'(1) = (3 - 2 \cdot 1)(3 - 1 - 1^2) + (3 \cdot 1 - 1^2)(-1 - 2 \cdot 1) = 1 + 2(-3) = -5$$

$$x_0 = 1, \quad y_0 = y(1) = (3 \cdot 1 - 1^2)(3 - 1 - 1^2) = 2 \cdot 1 = 2$$

$$y - 2 = -5(x - 1) \quad \square$$

3. Use the quotient rule and trigonometric identities to show...

3a. $\frac{d}{dx}[\tan(x)] = \sec^2(x)$

Sol:
$$\begin{aligned}\frac{d}{dx}[\tan(x)] &= \frac{d}{dx}\left[\frac{\sin(x)}{\cos(x)}\right] = \frac{\cos(x)\frac{d}{dx}[\sin(x)] - \sin(x)\frac{d}{dx}[\cos(x)]}{(\cos(x))^2} \\ &= \frac{\cos(x) \cdot \cos(x) - \sin(x) \cdot (-\sin(x))}{\cos^2(x)} \\ &= \frac{\cos^2(x) + \sin^2(x)}{\cos^2(x)} = \frac{1}{\cos^2(x)} = \sec^2(x) \quad \square\end{aligned}$$

3b. $\frac{d}{dx}[\sec(x)] = \sec(x)\tan(x)$

Sol:
$$\begin{aligned}\frac{d}{dx}[\sec(x)] &= \frac{d}{dx}\left[\frac{1}{\cos(x)}\right] = \frac{\cos(x)\frac{d}{dx}[1] - 1 \cdot \frac{d}{dx}[\cos(x)]}{(\cos(x))^2} \\ &= \frac{\cos(x) \cdot 0 - (-\sin(x))}{\cos^2(x)} \\ &= \frac{\sin(x)}{\cos^2(x)} = \frac{1}{\cos(x)} \cdot \frac{\sin(x)}{\cos(x)} = \sec(x)\tan(x) \quad \square\end{aligned}$$

3c. $\frac{d}{dx}[\csc(x)] = -\csc(x)\cot(x)$

Sol:
$$\begin{aligned}\frac{d}{dx}[\csc(x)] &= \frac{d}{dx}\left[\frac{1}{\sin(x)}\right] = \frac{\sin(x)\frac{d}{dx}[1] - 1 \cdot \frac{d}{dx}[\sin(x)]}{(\sin(x))^2} \\ &= \frac{\sin(x) \cdot 0 - \cos(x)}{\sin^2(x)} \\ &= -\frac{\cos(x)}{\sin^2(x)} = -\frac{1}{\sin(x)} \cdot \frac{\cos(x)}{\sin(x)} = -\csc(x)\cot(x) \quad \square\end{aligned}$$

3d. $\frac{d}{dx}[\cot(x)] = -\csc^2(x)$

Sol:
$$\begin{aligned}\frac{d}{dx}[\cot(x)] &= \frac{d}{dx}\left[\frac{\cos(x)}{\sin(x)}\right] = \frac{\sin(x)\frac{d}{dx}[\cos(x)] - \cos(x)\frac{d}{dx}[\sin(x)]}{(\sin(x))^2} \\ &= \frac{\sin(x) \cdot (-\sin(x)) - \cos(x) \cdot \cos(x)}{\sin^2(x)} \\ &= -\left(\frac{\sin^2(x) + \cos^2(x)}{\sin^2(x)}\right) \\ &= -\frac{1}{\sin^2(x)} = -\csc^2(x) \quad \square\end{aligned}$$

4. Calculate the indicated higher derivative.

4a. $\frac{d^2}{dx^2} \left[\frac{\cos(x)}{x} \right]$ $y = \frac{\cos(x)}{x}$

Sol: $\frac{dy}{dx} = \frac{d}{dx} \left[\frac{\cos(x)}{x} \right] = \frac{x \frac{d}{dx} [\cos(x)] - \cos(x) \frac{d}{dx} [x]}{x^2} = \frac{-x \sin(x) - \cos(x)}{x^2}$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} \left[\frac{-x \sin(x) - \cos(x)}{x^2} \right] = \frac{x^2 \frac{d}{dx} [-x \sin(x) - \cos(x)] - (-x \sin(x) - \cos(x)) \frac{d}{dx} [x^2]}{(x^2)^2}$$

$$= \frac{x^2 \left(\frac{d}{dx} [-x] \sin(x) + (-x) \frac{d}{dx} [\sin(x)] - (-\sin(x)) \right) + (-x \sin(x) - \cos(x)) (2x)}{x^4}$$

$$= \frac{x^2 (-\sin(x) - x \cos(x) - \cos(x)) + x (2x \sin(x) + 2 \cos(x))}{x^4}$$

$$= \frac{x (-x \sin(x) - x^2 \cos(x) - x \cos(x) + 2x \sin(x) + 2 \cos(x))}{x^4}$$

$$= \frac{(-x^2 - x + 2) \cos(x) + x \sin(x)}{x^3} \quad \square$$

4b. $\frac{d^3}{dx^3} [x^2 + 3x - 7]$ $y = x^2 + 3x - 7$

Sol: $\frac{dy}{dx} = 2x + 3$, $\frac{d^2y}{dx^2} = 2$, $\frac{d^3y}{dx^3} = 0$ \square

4c. $\frac{d^2}{dx^2} [\sec(x)]$ $y = \sec(x)$

Sol: $\frac{dy}{dx} = \frac{d}{dx} [\sec(x)] = \sec(x) \tan(x)$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} [\sec(x) \tan(x)] = \frac{d}{dx} [\sec(x)] \tan(x) + \sec(x) \frac{d}{dx} [\tan(x)]$$

$$= \sec(x) \tan(x) \cdot \tan(x) + \sec(x) \cdot \sec^2(x) = \sec(x) (\tan^2(x) + \sec^2(x))$$

4d. $\frac{d^2}{dx^2} [x \sin(x)]$ $y = x \sin(x)$

Sol: $\frac{dy}{dx} = \frac{d}{dx} [x] \sin(x) + x \frac{d}{dx} [\sin(x)] = \sin(x) + x \cos(x)$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} [\sin(x) + x \cos(x)] = \cos(x) + \frac{d}{dx} [x] \cos(x) + x \frac{d}{dx} [\cos(x)]$$

$$= 2 \cos(x) - x \sin(x) \quad \square$$